

PDES II, Math 8142 (old 562)

Prof. Gutiérrez Homework 3, Sobolev spaces (Due on 3/26/09)

Let $\Omega \subset \mathbb{R}^n$ be an open set, $k \geq 0$ an integer and $1 \leq p \leq \infty$. If $f, g \in L^1_{loc}(\Omega)$ and $\alpha = (\alpha_1, \dots, \alpha_n)$ is a multi-index we say that g is the α -th weak derivative of f if

$$\int_{\Omega} g(x)\phi(x) dx = (-1)^{|\alpha|} \int_{\Omega} f(x)D^{\alpha}\phi(x) dx,$$

and we write $g = D^{\alpha}f$. The Sobolev spaces are defined as follows:

$W^{k,p}(\Omega) = \{f \in L^p(\Omega) : \text{the } \alpha\text{-th weak derivative of } f, D^{\alpha}f \in L^p(\Omega) \text{ for all } |\alpha| \leq k\}$,

$$\|f\|_{k,p,\Omega} = \left(\sum_{|\alpha| \leq k} \|D^{\alpha}f\|_{L^p(\Omega)}^p \right)^{1/p} \approx \sum_{|\alpha| \leq k} \|D^{\alpha}f\|_{L^p(\Omega)}, \quad \text{if } p < \infty,$$

and

$$\|f\|_{k,\infty,\Omega} = \sum_{|\alpha| \leq k} \|D^{\alpha}f\|_{L^{\infty}(\Omega)}.$$

We also have the spaces

$$H^{k,p}(\Omega) = \overline{\{f \in C^{\infty}(\Omega) : \|f\|_{k,p,\Omega} < \infty\}};$$

$$W_0^{k,p}(\Omega) = \overline{\{f \in C_0^{\infty}(\Omega)\}},$$

where the closures are taken with respect to the norm $\|\cdot\|_{k,p,\Omega}$. We have

$$W_0^{k,p}(\Omega) \subset H^{k,p}(\Omega) \subset W^{k,p}(\Omega) \subset L^p(\Omega).$$

If $p < \infty$ then given $f \in W^{k,p}(\Omega)$ there exists a sequence $f_m \in C^{\infty}(\Omega) \cap W^{k,p}(\Omega)$ such that $f_m \rightarrow f$ in the norm $\|\cdot\|_{k,p,\Omega}$. That is

$$H^{k,p}(\Omega) = W^{k,p}(\Omega), \quad \text{if } 1 \leq p < \infty.$$

If Ω is bounded and has regular boundary, e.g., $\partial\Omega$ is C^1 , and $1 \leq p < \infty$, then

(a) Given $f \in W^{k,p}(\Omega)$ there exists a sequence $f_m \in C^{\infty}(\overline{\Omega})$ such that $f_m \rightarrow f$ in the norm $\|\cdot\|_{k,p,\Omega}$.

(b) There exists a bounded linear operator $T : W^{1,p}(\Omega) \rightarrow L^p(\partial\Omega)$ such that $\|Tf\|_{L^p(\partial\Omega)} \leq C\|f\|_{1,p,\Omega}$, with a constant $C > 0$ independent of f , and $Tf = f|_{\partial\Omega}$ for $f \in W^{1,p}(\Omega) \cap C(\overline{\Omega})$. Tf is the trace of f on $\partial\Omega$.

1. If f is continuous in Ω and the weak derivative $D^i f$ is also continuous in Ω , then the ordinary derivative f_{x_i} exists and is continuous in Ω .

HINT: enough to prove it in dimension one, that is, if

$$\int_a^b f(x)g'(x) dx = - \int_a^b h(x)g(x) dx \quad (1)$$

with f and h continuous and for all $g \in C_0^\infty(a, b)$, then $f'(x) = h(x)$ in (a, b) . Equation (1) is equivalent to

$$\int_a^b \left(f(x) - \int_{x_0}^x h(t) dt \right) g'(x) dx = 0,$$

so it is enough to show that if

$$\int_a^b w(x) g'(x) dx = 0, \quad \forall g \in C_0^\infty(a, b) \quad (2)$$

with w continuous, then w is constant in (a, b) . Now, if (2) holds then prove that $\int_a^b w(x) g'(x) dx = 0$ for all g absolutely continuous with compact support in (a, b) (take ϕ smooth with compact support in $[-1, 1]$, $\phi_\epsilon(x) = \epsilon^{-1} \phi(x/\epsilon)$ and take $\phi_\epsilon \star g$). Then given $a < a' < b' < b$, define $g(x) = \int_{a'}^x \left(w(t) - \int_{a'}^{b'} w(s) ds \right) dt$, for $a' \leq x \leq b'$, and $g(x) = 0$ outside $[a', b']$. Show that g is absolutely continuous. Notice that (2) holds if and only if $\int_a^b (w(x) - c) g'(x) dx = 0$ with any constant c . Then from this get that $\int_{a'}^{b'} \left(w(x) - \int_{a'}^{b'} w(s) ds \right)^2 dx = 0$.

2. Let $f(x) = |x_1|$ in the unit ball $B_1(0) \subset \mathbb{R}^n$. Show that f has weak derivatives of order α for all $|\alpha| \leq 1$ and $D^1 f = \text{sign } x_1$ and $D^i f = 0$ for $i = 2, \dots, n$.
3. Let $f \in L_{loc}^1(\Omega)$, Ω connected and $D^\alpha f = 0$ for all $|\alpha| = 1$. Then f is constant in Ω .
4. Assume Ω is regular. Prove that the trace of a function is unique.
5. Let Ω be bounded with C^1 boundary. Let $f, g \in W^{1,2}(\Omega)$. Prove the formula of integration by parts

$$\int_{\Omega} D^i f(x) g(x) dx = \int_{\partial\Omega} f(x) g(x) \eta_i d\sigma(x) - \int_{\Omega} f(x) D^i g(x) dx,$$

where η_i is the i -th component of the unit outer normal to $\partial\Omega$, and in the surface integral f and g are understood as the traces of f and g .

6. Let x_1, \dots, x_{k+1} be numbers and consider the Vandermonde determinant

$$V(x_1, \dots, x_{k+1}) = \det \begin{bmatrix} 1 & 1 & \dots & 1 \\ x_1 & x_2 & \dots & x_{k+1} \\ x_1^2 & x_2^2 & \dots & x_{k+1}^2 \\ \vdots & \vdots & \ddots & \vdots \\ x_1^k & x_2^k & \dots & x_{k+1}^k \end{bmatrix}.$$

We have that $V(x_1, \dots, x_{k+1}) = \prod_{1 \leq i < m \leq k+1} (x_m - x_i)$.

Let $B = B_1(0)$ be the unit ball in \mathbb{R}^n and $B^+ = \{x \in B_1(0) : x_n > 0\}$. Let $f \in C^k(\overline{B^+})$, write $x = (x', x_n)$ and define the extension

$$\bar{f}(x) = \begin{cases} f(x), & \text{for } x_n \geq 0 \\ \sum_{i=1}^{k+1} a_i f\left(x', -\frac{x_n}{i}\right), & \text{for } x_n < 0, \end{cases}$$

where $a_i, i = 1, \dots, k+1$, is the unique solution of the linear system

$$\sum_{i=1}^{k+1} \left(-\frac{1}{i}\right)^s a_i = 1, \quad s = 0, \dots, k.$$

Prove that $\bar{f} \in C^k(B)$ and

$$\|\bar{f}\|_{W^{k,p}(B)} \leq C \|f\|_{W^{k,p}(B^+)}.$$

7. Let $\lambda > 0$ and B the unit ball in \mathbb{R}^n . Show that there exists a constant $C > 0$ depending only on n and λ such that if $|\{x \in B : u(x) = 0\}| \geq \lambda$ then

$$\int_B u(x)^2 dx \leq C \int_B |\nabla u(x)|^2 dx.$$

8. Let $f \in C^1(\mathbb{R})$, $f' \in L^\infty(\mathbb{R})$ and $u \in W^{1,p}(\Omega)$. If $v(x) = f(u(x))$ then $v \in W^{1,p}(\Omega)$ and $Dv = f'(u)Du$.
9. Let $u \in W^{1,p}(\Omega)$ prove that

- (a) $u^+, u^- \in W^{1,p}(\Omega)$, and

$$\nabla u^+ = \begin{cases} \nabla u, & \text{a.e. on } \{u > 0\} \\ 0, & \text{a.e. on } \{u \leq 0\}, \end{cases}$$

$$\nabla u^- = \begin{cases} 0, & \text{a.e. on } \{u \geq 0\} \\ \nabla u, & \text{a.e. on } \{u < 0\}, \end{cases}$$

Hint: consider

$$h_\epsilon(x) = \begin{cases} (x^2 + \epsilon^2)^{1/2} - \epsilon, & \text{if } x \geq 0 \\ 0, & \text{if } x < 0, \end{cases}$$

and show that $u^+ = \lim_{\epsilon \rightarrow 0} h_\epsilon(u)$.

- (b) $\nabla u = 0$ a.e. on $\{u = 0\}$.
- (c) $|u| \in W^{1,p}(\Omega)$.