



ELSEVIER

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

SCIENCE @ DIRECT®

LINEAR ALGEBRA  
AND ITS  
APPLICATIONS

Linear Algebra and its Applications 363 (2003) 65–80

[www.elsevier.com/locate/laa](http://www.elsevier.com/locate/laa)

## Conditions for strict inequality in comparisons of spectral radii of splittings of different matrices

Ludwig Elsner <sup>a</sup>, Andreas Frommer <sup>b,\*</sup>, Reinhard Nabben <sup>a</sup>,  
Hans Schneider <sup>c,1</sup>, Daniel B. Szyld <sup>d,2</sup>

<sup>a</sup>Fakultät für Mathematik, Universität Bielefeld, Postfach 10 01 31, D-33501 Bielefeld, Germany

<sup>b</sup>Fachbereich Mathematik, Universität Wuppertal, D-42097 Wuppertal, Germany

<sup>c</sup>Department of Mathematics, University of Wisconsin at Madison, 480 Lincoln Drive,  
Madison, WI 53706, USA

<sup>d</sup>Department of Mathematics, Temple University, Philadelphia, PA 19122-6094, USA

Received 16 March 2001; accepted 4 October 2001

Submitted by D. Hershkowitz

---

### Abstract

We present new comparison theorems for the spectral radii of matrices arising from splittings of different matrices under nonnegativity assumptions. Our focus is on establishing strict inequalities of the spectral radii without imposing strict inequalities of the matrices, but we also obtain new results for nonstrict inequalities of the spectral radii. We emphasize two different approaches, one combinatorial and the other analytic and discuss their merits in the light of the results obtained. We try to get fairly general results and indicate by counter-examples that some of our hypotheses cannot be relaxed in certain directions.

© 2001 Elsevier Science Inc. All rights reserved.

*AMS classification:* 65F10; 15A06; 15A48

*Keywords:* Strict comparison theorems; Combinatorial matrix theory; Nonnegative matrices; Perron–Frobenius theory; Iterative solution of linear systems

---

\* Corresponding author.

*E-mail addresses:* [elsner@mathematik.uni-bielefeld.de](mailto:elsner@mathematik.uni-bielefeld.de) (L. Elsner), [frommer@math.uni-wuppertal.de](mailto:frommer@math.uni-wuppertal.de) (A. Frommer), [nabben@mathematik.uni-bielefeld.de](mailto:nabben@mathematik.uni-bielefeld.de) (R. Nabben), [hans@math.wisc.edu](mailto:hans@math.wisc.edu) (H. Schneider), [szyld@math.temple.edu](mailto:szyld@math.temple.edu) (D.B. Szyld).

<sup>1</sup> Research supported by the US Social Security Administration and the Wisconsin Department of Employee Trust Funds.

<sup>2</sup> Supported by the National Science Foundation grant DMS-9973219.

## 1. Introduction

We study comparison theorems between nonnegative splittings of two different matrices. Our focus is on strict inequalities for the spectral radii, while the inequalities in the hypotheses (with respect to the nonnegative cone of matrices) are not strict.

We present two fundamentally different approaches to this topic. The first one is combinatorial in nature and makes explicit use of the graph of the matrix involved. The second approach is analytic in nature and relies on topological and algebraic arguments. Both these approaches are interesting by their methodology. They usually complement each other. For example, in [13,14] graph theoretical arguments were used to prove certain results, while in [9] analytical arguments were used for the same results. In our comparison theorems here it turns out that the graph theoretical approach is somehow restricted to considering  $M$ -matrices, whereas the analytical approach allows us to obtain further results involving general monotone matrices, i.e., matrices with nonnegative inverses.

Comparison theorems between the spectral radii of matrices are a useful tool in the analysis of the rate of convergence of iterative methods or for judging the efficiency of preconditioners. There is also a connection to population dynamics; see, e.g., [7] and the references given therein.

The paper is organized as follows. In Section 2 we derive a strict inequality result using the combinatorial approach. We give two different versions of this result. The first applies to iterative methods obtained through matrix splittings, whereas the second is more appropriate, e.g., when studying population dynamics. In Section 3 we then present several generalizations of this result in the splitting formulation using an analytic approach. There, we introduce and use various notions of nonnegative splittings. Further extensions are presented in Section 4. In particular, one can appreciate the variety of comparison results that can be obtained for splittings of different matrices. This section also contains a new result for nonstrict inequalities.

## 2. The combinatorial approach

We use the notation  $A \geq A'$  for two real matrices of equal size if each entry of the difference  $A - A'$  is nonnegative. We write  $A > A'$  if each entry of the difference is positive. A matrix  $A \geq O$  ( $A > O$ ) is called nonnegative (positive). We will often consider relations  $A \geq A'$  with equality excluded ( $A \neq A'$ ) for which we write  $A \geq A'$ .

A nonsingular  $M$ -matrix  $A$  is such that it can be written as  $A = \sigma I - T$  with  $T \geq O$  and  $\sigma > \rho(T)$ ; see, e.g., [2,15]. Alternatively,  $A$  is a nonsingular  $M$ -matrix if it can be expressed as  $A = \sigma(I - T')$  with  $\sigma > 0$ ,  $T' \geq O$  and  $\rho(T') < 1$ .

Given a square matrix  $A$ ,  $A = M - N$  is called a splitting if  $M$  is nonsingular. A splitting is *regular* if  $M^{-1} \geq O$  and  $N \geq O$  [15]. It is an  *$M$ -splitting* if  $M$  is an  $M$ -matrix and  $N \geq O$  [13]; see further Definition 3.3.

We can state now our main theorem for splittings of  $M$ -matrices.

**Theorem 2.1.** Let  $A_1 = M_1 - N$  and  $A_2 = M_2 - N$  be  $M$ -splittings of  $A_1$  and  $A_2$ , respectively, where

$$M_1 \preceq M_2 \quad \text{and} \quad N \succeq O. \quad (1)$$

If  $A_1^{-1} > O$ , then

$$0 < \rho(NM_2^{-1}) < \rho(NM_1^{-1}). \quad (2)$$

The importance of this theorem is that (2) is strict while hypotheses (1) are not. As we shall see, Theorem 2.1 can be considered a reformulation of part (2) of the following result.

**Theorem 2.2.** (i) Let  $T$  and  $F$  be square nonnegative matrices. Suppose that the spectral radius satisfies  $\rho(T) < 1$ . Assume also that  $F \neq O$  and that  $F + T$  is irreducible. Let  $Q = F(I - T)^{-1}$ . Then, after a permutation similarity,

$$Q = \begin{pmatrix} Q_{11} & Q_{12} \\ O & O \end{pmatrix},$$

where  $Q_{11}$  is a nontrivial irreducible nonnegative matrix,  $Q_{12}$  is a nonnegative matrix every column of which has a positive entry, and the zero rows of  $Q$  correspond to the zero rows of  $F$ , if any. Further  $\rho(Q) > 0$ .

(ii) In addition, let  $O \leq T' \preceq T$ , and put  $Q' = F(I - T')^{-1}$ . Then, after the same permutation similarity and partitioning,

$$Q' = \begin{pmatrix} Q'_{11} & Q'_{12} \\ O & O \end{pmatrix},$$

where  $Q'_{11} \preceq Q_{11}$  and  $\rho(Q') < \rho(Q)$ .

Again, we observe that the point of the theorem is that the inequalities on the spectral radii in Parts (i) and (ii) are *strict*, since it is trivial that  $O \leq Q' \leq Q$  and hence it follows by standard Perron–Frobenius theory that  $0 \leq \rho(Q') \leq \rho(Q)$ .

We will present a complete graph theoretic proof of Theorem 2.2, although Part (i) was presented, in terms of the  $M$ -splitting of Theorem 2.1, at the Linear Algebra meeting Oberwolfach in 1982 and appeared as Lemma 3.4 of [13]; see also [14,17].

Before starting the proof, let us introduce standard terminology for graphs of nonnegative matrices; see, e.g., [13]. Let  $n$  be a positive integer. Then a *path* (without further qualification) will be a sequence  $p = (i(0), \dots, i(s))$  of positive integers  $i(r)$ ,  $1 \leq i(r) \leq n$ ,  $r = 0, \dots, s$ . A path is usually called an *arc* if  $s = 1$ . Let  $A$  be a nonnegative  $n \times n$  matrix. We call the path  $p$  an *A-path* or a *path in A* if  $a_{i(r-1), i(r)} > 0$ ,  $r = 1, \dots, s$ , and we use similar terminology for arcs.

For any path  $p$ , the *path product*  $p(A)$  is defined by

$$p(A) = a_{i(0),i(1)} \cdots a_{i(s-1),i(s)}.$$

Let  $F$  and  $T$  be two nonnegative  $n \times n$  matrices and suppose the spectral radius of  $T$  satisfies  $\rho(T) < 1$ . We are here concerned with the matrix  $Q = F(I - T)^{-1}$ , as essentially was [13] where  $Q^T$  was considered. Since  $Q = F(I + T + T^2 + \cdots)$ , the elements of  $Q$  are easily computed to be

$$q_{i,j} = \sum_{\alpha, p} \alpha(F)p(T), \tag{3}$$

where the summation is taken over all arcs  $\alpha = (i, k)$  and all paths  $p$  from  $k$  to  $j$  or—considering only nonzero summands—over all  $F$ -arcs  $\alpha = (i, k)$  and all  $T$ -paths  $p$  from  $k$  to  $j$ , see [13, Theorem 2.7]. In view of this, we call a path *relevant* if its first arc is an  $F$ -arc and all other arcs (if any) are  $T$ -arcs. By (3),  $q_{i,j} > 0$  if and only if there is a relevant path from  $i$  to  $j$ . If  $q = (\alpha, p)$  is a relevant path with first arc  $\alpha$ , we define  $q(F, T) = \alpha(F)p(T)$ . Hence (3) may be rewritten as

$$q_{i,j} = \sum_{q \in \mathfrak{R}(i,j)} q(F, T), \tag{4}$$

where  $\mathfrak{R}(i, j)$  is the set of all relevant paths from  $i$  to  $j$ .

Our proofs depend on the following graph theoretic remark:

**Remark 2.3.** Let  $p$  be an  $(F + T)$ -path that begins with an  $F$ -arc. Then  $p$  may be decomposed as  $p = (q_1, \dots, q_s)$ , where each  $q_\ell, \ell = 1, \dots, s$ , is a relevant path. As an additional condition, we may even impose that each  $q_\ell, \ell = 2, \dots, s$ , begins with an  $F$ -arc which is *not* a  $T$ -arc. Since each  $q_\ell$  begins with an  $F$ -arc which corresponds to a positive element of  $Q$ , it follows that, if  $p$  is an  $(F + T)$ -path from  $i$  to  $j$  that begins with an  $F$ -arc, there is a  $Q$ -path from  $i$  to  $j$  (and the converse also holds).

**Proof of Theorem 2.2.** Let  $\mathcal{B}$  be the set of vertices which are the starts of  $F$ -arcs and let  $\mathcal{A}$  be the complement of  $\mathcal{B}$  in  $\{1, \dots, n\}$ . Apply a permutation similarity to all matrices so that  $\mathcal{B} = \{1, \dots, k\}$ ,  $1 \leq k \leq n$ . Note that  $\mathcal{B}$  is nonempty and partition  $Q$  so that  $Q_{11}$  is  $k \times k$ .

Let  $b$  be any vertex of  $\mathcal{B}$  and let  $d$  be any vertex. Let  $\alpha = (b, c)$  be an  $F$ -arc. Since  $F + T$  is irreducible, there exists an  $(F + T)$ -path  $r$  from  $c$  to  $d$  and let  $p = (\alpha, r)$ . Suppose that  $p = (q_1, \dots, q_s)$  is a decomposition into relevant paths. In view of Remark 2.3 it follows that there is a  $Q$ -path from  $b$  to  $d$ . Hence there is a nonempty path in  $Q$  from every vertex of  $\mathcal{B}$  to every other vertex. It follows that every column of  $[Q_{11} \ Q_{12}]$  contains a positive element. Now let  $d \in \mathcal{B}$ . Then every  $q_\ell, \ell = 1, \dots, s$ , corresponds to a relevant path beginning and ending at an element of  $Q_{11}$ . Hence there exists a  $Q_{11}$ -path from  $b$  to  $d$  and it follows by a standard result that  $Q_{11}$  is irreducible. Also, there is no relevant path from a vertex of  $\mathcal{A}$  since there is no  $F$ -arc that starts there. This proves that  $Q_{21} = O$  and  $Q_{22} = O$ . Since  $Q_{11}$  is a nontrivial irreducible matrix (viz.  $Q_{11} \neq O$ ), we have  $0 < \rho(Q_{11}) = \rho(Q)$ . This proves Part (i).

Since  $O \leq T' \leq T$ , there exists an arc  $(i, j)$  of  $T$  such that  $0 \leq t'_{i,j} < t_{i,j}$ . Let  $b$  be any vertex of  $\mathcal{B}$  and let  $(b, c)$  be an  $F$ -arc. Since  $F + T$  is irreducible, there exists an  $(F + T)$ -path  $p$  which begins with the arc  $(b, c)$ , then continues through the arc  $(i, j)$  (repeating  $(b, c)$  if it happens that  $(b, c) = (i, j)$ ) and continues back to  $b$ . We decompose the path  $p = (q_1, \dots, q_s)$  into relevant paths that satisfy the additional condition that each  $q_\ell, \ell > 1$ , begins with an  $F$ -arc which is not a  $T$ -arc. Then there is an  $\ell, 1 \leq \ell \leq s$ , such that  $(i, j)$  is a noninitial arc of  $q_\ell$  and  $q_\ell(F, T') < q_\ell(F, T)$ , since  $t'_{i,j} < t_{i,j}$ . The relevant path  $q_\ell$  begins and ends at vertices of  $\mathcal{B}$ ; say it is an  $(a, d)$  path. Since for all relevant paths  $q$  we have  $q(F, T') \leq q(F, T)$ , it follows by (4) that  $q'_{a,d} < q_{a,d}$ . This proves that  $Q'_{11} \leq Q_{11}$  and, since  $Q_{11}$  is irreducible, it follows that  $\rho(Q') = \rho(Q'_{11}) < \rho(Q_{11}) = \rho(Q)$ , and Part (ii) is proved.  $\square$

**Remark 2.4.** Above we have essentially proved one direction of the following result:  $q'_{a,d} < q_{a,d}$  if and only if there exists a relevant path from  $a$  to  $d$  which contains a noninitial arc  $(i, j)$  such that  $t'_{i,j} < t_{i,j}$ .

As a corollary of Part (ii) of Theorem 2.2 we have the following result first proved as [7, Theorem 4.4].

**Proposition 2.5.** Let  $T$  and  $F$  be square nonnegative matrices with  $F \neq O$  and  $F + T$  irreducible. Suppose that the spectral radius  $\rho(T)$  of  $T$  satisfies  $\rho(T) < 1$ . Then  $\rho(F(I - T)^{-1}) > 0$ . Moreover, let  $s > \rho(T)$ . Then

$$\rho(F(I - T)^{-1}) < \rho\left(F\left(I - \frac{1}{s}T\right)^{-1}\right) \quad \text{if } s < 1$$

and

$$\rho(F(I - T)^{-1}) > \rho\left(F\left(I - \frac{1}{s}T\right)^{-1}\right) \quad \text{if } s > 1.$$

The above proposition has a direct interpretation in models from population dynamics where  $T$  and  $F$  represent the transition and fertility matrix, respectively, and  $s$  is the growth rate. We refer to [7] for details.

We now turn to prove Theorem 2.1 using Theorem 2.2.

**Proof of Theorem 2.1.** Since  $M_i, i = 1, 2$ , is an  $M$ -matrix, we may write  $M_i = c_i(I - \tilde{T}_i)$ , where  $c_i > 0, \tilde{T}_i \geq O, \rho(\tilde{T}_i) < 1$  for  $i = 1, 2$ . Let  $c = \max\{c_1, c_2\}$  and write

$$M_i = c_i(I - \tilde{T}_i) = c\left(I - \left(\frac{c - c_i}{c}I + \frac{c_i}{c}\tilde{T}_i\right)\right) = c(I - T_i).$$

Here,

$$T_i = \frac{c - c_i}{c}I + \frac{c_i}{c}\tilde{T}_i \geq O \quad \text{and} \quad \rho(T_i) = \frac{c - c_i}{c} + \frac{c_i}{c}\rho(\tilde{T}_i) < 1.$$

Since  $O < A_1^{-1} = c^{-1}(I - N - T_1)^{-1}$ , it follows that  $N + T_1$  is irreducible. Thus the theorem is implied by Theorem 2.2 (where  $F = N$ ,  $T = T_1$ ,  $T' = T_2$ ).  $\square$

The proof just presented shows that Theorem 2.2 implies Theorem 2.1. If we disregard the statement on the structure of  $Q$  and  $Q'$  in Theorem 2.2, the converse also holds: Theorem 2.1 implies Theorem 2.2. This follows indeed easily, since in Theorem 2.2 we may without loss of generality scale  $F = N$  by a positive constant so that  $\rho(F + T) < 1$ , which yields  $(I - F - T)^{-1} > O$ .

We end this section with a couple of examples showing that the hypotheses  $M_1 \preceq M_2$  and  $A_1^{-1} > O$  in Theorem 2.1 cannot be weakened.

First observe that for  $M$ -matrices  $M_1, M_2$ ,  $M_1 \preceq M_2$  implies that  $M_1^{-1} \succeq M_2^{-1}$  but not conversely. It is tempting to assume that Theorem 2.1 remains valid if the hypothesis  $M_1 \preceq M_2$  is replaced by  $M_1^{-1} \succeq M_2^{-1}$ , but this is not so as is easily shown by the following example.

**Example 2.6.** Let

$$M_1^{-1} = \begin{pmatrix} 1 & .5 \\ .5 & 1 \end{pmatrix}, \quad M_2^{-1} = \begin{pmatrix} 1 & .5 \\ .4 & 1 \end{pmatrix}, \quad N = \begin{pmatrix} .1 & 0 \\ 0 & 0 \end{pmatrix}.$$

Then  $M_1^{-1} \succeq M_2^{-1}$  and  $(M_1 - N)^{-1} > O$  but  $NM_1^{-1} = NM_2^{-1}$ .

**Remark 2.7.** If we assume *strict* inequality  $M_1^{-1} > M_2^{-1}$ , the other assumptions of Theorem 2.1 remaining unchanged, then the conclusion  $\rho(NM_2^{-1}) < \rho(NM_1^{-1})$  still holds. This can be seen by considering the reducible normal form; see, e.g., [15]. Observe that  $O \leq NM_2^{-1} \preceq NM_1^{-1}$  follows immediately from the assumptions, and  $NM_1^{-1}$  has at least one positive diagonal element. So the reducible normal form of  $NM_1^{-1}$  has at least one nontrivial (irreducible) diagonal block. But all positive elements in each such block are strictly decreased when passing from  $NM_1^{-1}$  to  $NM_2^{-1}$ , so the spectral radius of each block strictly decreases. This gives  $\rho(NM_1^{-1}) > \rho(NM_2^{-1})$  by standard Perron–Frobenius theory.

It is also easy to see that we cannot omit the condition that  $A_1^{-1} > O$  from Theorem 2.1 (or, equivalently, that  $F + T$  is irreducible in Theorem 2.2).

**Example 2.8.** Let

$$N = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}, \quad M_1 = \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix}, \quad M_2 = I.$$

Then

$$NM_1^{-1} = NM_2^{-1} = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}.$$

### 3. An analytic approach

The purpose of this section is to develop generalizations for Theorem 2.1. We will be able to dispense with the  $M$ -splitting hypothesis by assuming more general non-negativity hypotheses. Section 4 will contain further generalizations together with a discussion of possible applications; see Remark 4.7. Our approach is now analytical rather than combinatorial as exemplified by the following auxiliary result.

**Lemma 3.1.** *Let  $B_0$  and  $B_1$  be two square nonsingular matrices such that  $B_0 \preceq B_1$  and  $B_0^{-1} > O$  as well as  $B_1^{-1} \geq O$ . Then  $B_0^{-1} > B_1^{-1}$ .*

**Proof.** Define the family of matrices  $B_\alpha = (1 - \alpha)B_0 + \alpha B_1$ ,  $\alpha \in [0, 1]$ . Clearly, we have  $B_\alpha \preceq B_\beta$  for  $\alpha < \beta$ . If  $B_\alpha, B_\beta$  are nonsingular, we also have

$$B_\alpha^{-1} - B_\beta^{-1} = B_\beta^{-1} (B_\beta - B_\alpha) B_\alpha^{-1}. \quad (5)$$

By continuity, there exists a small positive  $\gamma \in (0, 1)$  such that  $B_\gamma$  is nonsingular with  $B_\gamma^{-1} > O$ . Since  $B_\gamma - B_0$  is nonnegative with at least one positive entry, the product  $B_\gamma^{-1}(B_\gamma - B_0)$  has at least one positive column. Since  $B_0^{-1} > O$ , the product  $B_\gamma^{-1}(B_\gamma - B_0)B_0^{-1}$  is therefore positive. From (5) (with  $\alpha = 0, \beta = \gamma$ ) we therefore get

$$B_0^{-1} > B_\gamma^{-1}. \quad (6)$$

We also have  $B_1 - B_\gamma \geq O$ , and since  $B_1^{-1} \geq O, B_\gamma^{-1} > O$  we have

$$B_\gamma^{-1} \geq B_1^{-1}, \quad (7)$$

again from (5) (with  $\alpha = \gamma, \beta = 1$ ). Combining (6) and (7) we obtain the desired strict inequality  $B_0^{-1} > B_1^{-1}$ .  $\square$

We also need the following, fairly standard auxiliary result; see, e.g., [8]. For the sake of completeness, we reproduce a proof here. Note that  $T$  need not be irreducible.

**Lemma 3.2.** *Let  $T$  be a nonnegative matrix, let  $x$  be a nonnegative nonzero vector and  $\alpha$  a positive scalar.*

- (i) *If  $Tx \geq \alpha x$ , then  $\rho(T) \geq \alpha$ . Moreover, if  $Tx > \alpha x$ , then  $\rho(T) > \alpha$ .*
- (ii) *If  $Tx < \alpha x$ , then  $\rho(T) < \alpha$ .*

**Proof.** To show (i), assume that  $Tx \geq \alpha x$  but  $\rho(T) < \alpha$ . Then

$$\left(I - \frac{1}{\alpha}T\right)^{-1} = \sum_{\nu=0}^{\infty} \left(\frac{1}{\alpha}T\right)^{\nu} \geq O.$$

Therefore,

$$0 \leq \left(I - \frac{1}{\alpha}T\right)^{-1} (Tx - \alpha x) = -\alpha x.$$

Since we had  $x \geq 0$  this implies  $x = 0$ , a contradiction. This proves Part (i) for the nonstrict inequalities. If  $Tx > \alpha x$  we can choose  $\alpha_1 > \alpha$  for which still  $Tx \geq \alpha_1 x$ . By what we have already proved we get  $\rho(T) \geq \alpha_1$  which implies  $\rho(T) > \alpha$ .

To show (ii), note that  $x > 0$ . The matrix  $\frac{1}{\alpha}T$  then satisfies  $\frac{1}{\alpha}Tx < \beta x$  with  $0 < \beta < 1$ . This implies

$$0 \leq \left(\frac{1}{\alpha}T\right)^n x < \beta^n x, \quad n = 0, 1, \dots$$

This shows that  $\left(\frac{1}{\alpha}T\right)^n$  tends to zero as every entry tends to zero, so that  $\rho(T) < \alpha$ .  $\square$

In order to formulate and appreciate the generalization of Theorem 2.1 let us recall the following—now near-standard—terminology; see, e.g., [3,16].

**Definition 3.3.** A splitting  $A = M - N$  is called:

- (i) *weak nonnegative of first type* if  $M^{-1} \geq O$  and  $M^{-1}N \geq O$ ,
- (ii) *weak nonnegative of second type* if  $M^{-1} \geq O$  and  $NM^{-1} \geq O$ .
- (iii) *nonnegative* if it is weak nonnegative of both types.

Clearly, a regular splitting (and thus an  $M$ -splitting) is also weak nonnegative, and a weak nonnegative splitting is weak nonnegative of either type.

The following known results on weak nonnegative splittings are very important for our investigations. As it is usually done, we state these results in terms of the spectral radius of the “iteration matrix”  $M^{-1}N$ . But note that we could as well take  $NM^{-1}$  (as we did in Theorem 2.1) since  $\rho(M^{-1}N) = \rho(NM^{-1})$ .

**Theorem 3.4.** Let  $A$  be a nonsingular matrix with  $A^{-1} \geq O$ .

- (i) Let  $A = M - N$  be a weak nonnegative splitting of either type. Then  $\rho(M^{-1}N) < 1$ .
- (ii) If the splitting  $A = M - N$  is weak nonnegative of second type, there exists a vector  $x \geq 0$  such that  $M^{-1}Nx = \rho(M^{-1}N)x$  and  $Ax \geq 0$  as well as  $Nx \geq 0$ .

**Proof.** For the first type, Part (i) goes back to [11], whereas for second type splittings it was given in [16]. The major Part of (ii) was proved in [1, Lemma 2.8] except for the inequality  $Nx \geq 0$ . To prove this, let us write  $\rho = \rho(M^{-1}N)$ . Note that we have  $Nx = \rho Mx$  which gives  $Mx = (1/\rho)Nx$ . Therefore, we obtain

$$0 \leq Ax = M(I - M^{-1}N)x = (1 - \rho)Mx = \frac{1 - \rho}{\rho}Nx$$

with a positive factor  $(1 - \rho)/\rho$  since  $\rho < 1$ . So  $Nx \geq 0$ , and equality is excluded because otherwise  $Ax = 0$  with  $x \neq 0$  which is impossible.  $\square$

We now turn to the announced generalization of Theorem 2.1.

**Theorem 3.5.** Assume that  $A_1 = M_1 - N$ ,  $A_2 = M_2 - N$  are two weak nonnegative splittings of different types of nonsingular square matrices  $A_1, A_2$  with  $N \neq O$ . Assume that  $A_1 \preceq A_2$  (or, equivalently,  $M_1 \preceq M_2$ ) and that  $A_1^{-1} > O$ ,  $A_2^{-1} \geq O$ . Then  $\rho(M_2^{-1}N) < \rho(M_1^{-1}N) < 1$ .

**Proof.** By Theorem 3.4(i) we know that  $\rho(M_1^{-1}N) < 1$  and  $\rho(M_2^{-1}N) < 1$ . Denote  $G_1 = A_1^{-1}N$ ,  $G_2 = A_2^{-1}N$  and  $\tilde{G}_1 = NA_1^{-1}$ ,  $\tilde{G}_2 = NA_2^{-1}$ . We have

$$\begin{aligned} G_i &= A_i^{-1}N = (I - M_i^{-1}N)^{-1}(M_i^{-1}N), \quad i = 1, 2, \\ \tilde{G}_i &= NA_i^{-1} = NM_i^{-1}(I - NM_i^{-1})^{-1}, \quad i = 1, 2. \end{aligned}$$

Now first assume that  $A_1 = M_1 - N$  is of first type whereas  $A_2 = M_2 - N$  is of second type. We thus have  $M_1^{-1}N \geq O$  and  $NM_2^{-1} \geq O$  so that  $G_1$  and  $\tilde{G}_2$  are nonnegative matrices. Since  $\rho(G_i) = \rho(\tilde{G}_i)$  and since the function  $f : t \mapsto t/(1-t)$  is strictly increasing on  $[0, 1)$ , we have

$$\begin{aligned} \rho(G_i) &= \rho(\tilde{G}_i) = \rho(M_i^{-1}N)/(1 - \rho(M_i^{-1}N)) \\ &= \rho(NM_i^{-1})/(1 - \rho(NM_i^{-1})) \end{aligned}$$

for  $i = 1, 2$ . Thus all we need to show is  $\rho(G_2) < \rho(G_1)$ . By Theorem 3.4(ii) there exists a vector  $x \geq 0$  such that  $M_2^{-1}Nx = \rho(M_2^{-1}N)x$  and  $Nx \geq 0$ . By Lemma 3.1 we have  $A_1^{-1} > A_2^{-1}$ , so that together with  $A_1^{-1} > O$  we get

$$G_1x = A_1^{-1}Nx > A_2^{-1}Nx = G_2x = \rho(G_2)x. \quad (8)$$

Whence  $\rho(G_2) < \rho(G_1)$  by Lemma 3.2(i).

If  $NM_1^{-1} \geq O$  and  $M_2^{-1}N \geq O$  then  $\tilde{G}_1$  and  $G_2$  are nonnegative matrices. Again by Theorem 3.4(ii) there exists a nonzero nonnegative vector  $z$  such that  $M_1^{-1}Nz = \rho(M_1^{-1}N)z$  and  $Nz \geq 0$ . Thus

$$\rho(G_1)z = G_1z = A_1^{-1}Nz > A_2^{-1}Nz = G_2z, \quad (9)$$

and we obtain  $\rho(G_2) < \rho(G_1)$  by Lemma 3.2(ii).  $\square$

As a first comment, let us note that in the proof we only made use of the inequality  $A_1^{-1} - A_2^{-1} > O$ , but not of  $A_1 \preceq A_2$ . In the light of Lemma 3.1 a slightly more general version of the theorem therefore arises if one replaces the assumption  $A_1 \preceq A_2$  by  $A_1^{-1} - A_2^{-1} > O$ .

In the following corollaries we now emphasize two special cases of Theorem 3.5. In particular, these corollaries resemble Theorem 2.1 where the hypothesis of  $M$ -splittings has been replaced with the hypothesis of either regular or nonnegative splittings.

**Corollary 3.6.** Assume that  $A_1 = M_1 - N$ ,  $A_2 = M_2 - N$  are two regular splittings of square nonsingular matrices  $A_1, A_2$  with  $N \neq O$ . Moreover, let  $A_1 \preceq A_2$  (or, equivalently,  $M_1 \preceq M_2$ ) and assume that  $A_1^{-1} > O, A_2^{-1} \geq O$ . Then  $\rho(M_2^{-1}N) < \rho(M_1^{-1}N) < 1$ .

**Corollary 3.7.** Assume that  $A_1 = M_1 - N, A_2 = M_2 - N$  are two nonnegative splittings of square nonsingular matrices  $A_1, A_2$  with  $N \neq O$ . Moreover, let  $A_1 \preceq A_2$  (or, equivalently,  $M_1 \preceq M_2$ ) and assume that  $A_1^{-1} > O, A_2^{-1} \geq O$ . Then  $\rho(M_2^{-1}N) < \rho(M_1^{-1}N) < 1$ .

We comment now on how some of the hypotheses of the results in this section cannot be weakened.

For regular splittings, we cannot weaken the assumption  $A_1^{-1} > O$  by replacing it by the hypothesis that  $A_1^{-1}$  is irreducible, for in Example 3.8 below  $A_1^{-1}$  is irreducible and indeed has precisely one element equal to 0.

**Example 3.8.** Let

$$M_1 = \begin{pmatrix} 20 & -10 & -10 \\ -10 & 15 & -9 \\ -10 & 5 & 15 \end{pmatrix}, \quad M_2 = \begin{pmatrix} 20 & -10 & -10 \\ -10 & 15 & -6 \\ -10 & 5 & 15 \end{pmatrix},$$

$$N = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 2 \end{pmatrix},$$

and put  $A_1 = M_1 - N, A_2 = M_2 - N$ . Here  $M_2$  has just one positive off-diagonal element. Then, as we show explicitly below,  $M_2^{-1}, M_1^{-1}$ , and  $A_1^{-1}$  are nonnegative and only the (3, 2)-element is 0 in each case.

$$A_1^{-1} = \frac{1}{80} \begin{pmatrix} 12 & 4 & 12 \\ 11 & 8 & 14 \\ 5 & 0 & 10 \end{pmatrix},$$

$$M_1^{-1} = \frac{1}{200} \begin{pmatrix} 27 & 10 & 24 \\ 24 & 20 & 28 \\ 10 & 0 & 20 \end{pmatrix}, \quad M_2^{-1} = \frac{1}{400} \begin{pmatrix} 51 & 20 & 42 \\ 42 & 40 & 44 \\ 20 & 0 & 40 \end{pmatrix}.$$

But  $\rho(M_1^{-1}N) = \rho(M_2^{-1}N) = 1/5$ .

#### 4. Further results

The purpose of this section is to formulate additional comparison results for splittings  $A_1 = M_1 - N_1, A_2 = M_2 - N_2$  with two possibly different matrices  $N_1, N_2$ . Before we do so, we take a closer look at known results for two splittings of a *single* matrix  $A$  which we summarize in the following theorem.

**Theorem 4.1.** Assume that  $A$  is a nonsingular matrix such that  $A^{-1} > O$  and let  $A = M_1 - N_1$  and  $A = M_2 - N_2$  be two splittings of  $A$ . Then  $\rho(M_1^{-1}N_1) < \rho(M_2^{-1}N_2)$  in the following cases:

- (i) both splittings are regular,  $N_1 \neq 0$  and  $M_1 \preceq M_2$  or, equivalently,  $N_1 \preceq N_2$  [15, Theorem 3.32];
- (ii) both splittings are regular and  $M_1^{-1} > M_2^{-1}$  [(16, Theorem 3.6], see also [4]);
- (iii) both splittings are weak nonnegative splittings of different types and  $M_1^{-1} > M_2^{-1}$  [3, Theorem 7].

Clearly, Part (iii) contains the other two as well as an additional comparison result by Elsner [5] where one splitting is regular and the other is weak nonnegative of first type.

Note that Part (i) uses the same hypothesis  $M_1 \preceq M_2$  as we used in Theorem 3.5, but it establishes the reverse inequality between the spectral radii. This is no contradiction though, since in Theorem 3.5 the assumption  $A_1 \preceq A_2$  excludes equality between  $A_1$  and  $A_2$ .

Taking Theorem 4.1 as a source of inspiration, and as was done in [8,10], we will now formulate comparison theorems for two splittings  $A_1 = M_1 - N_1$ ,  $A_2 = M_2 - N_2$  of different matrices with similar hypotheses as in Parts (i) and (iii). As before, the emphasis is on strict inequality of the spectral radii, without always having strict inequalities in the hypotheses. As it turns out, requiring  $M_2 - M_1 \succeq O$  or  $M_1^{-1} - M_2^{-1} > O$  does not suffice. We need to bound these differences by  $A_2 - A_1$  and  $A_1^{-1} - A_2^{-1}$ , respectively.

**Theorem 4.2.** Assume that  $A_1 = M_1 - N_1$ ,  $A_2 = M_2 - N_2$  are two weak nonnegative splittings of different types of nonsingular matrices  $A_1, A_2$  with  $N_1, N_2 \neq O$ . Assume that  $A_1^{-1} - A_2^{-1} > O$  and that  $A_1^{-1} > O, A_2^{-1} \geq O$ .

(i) If

$$M_2 - M_1 \leq A_2 - A_1, \quad (10)$$

then

$$\rho(M_2^{-1}N_2) < \rho(M_1^{-1}N_1) < 1. \quad (11)$$

(ii) If  $\rho(M_1^{-1}N_1) > 0$  and

$$M_1^{-1} - M_2^{-1} \geq A_1^{-1} - A_2^{-1}, \quad (12)$$

then

$$\rho(M_1^{-1}N_1) < \rho(M_2^{-1}N_2) < 1. \quad (13)$$

**Proof.** By Theorem 3.4(i) we know that  $\rho_1 = \rho(M_1^{-1}N_1) < 1$ ,  $\rho_2 = \rho(M_2^{-1}N_2) < 1$ . So we only have to prove the first inequalities in (11) and (13). To prove Part (i)

note first that (10) is equivalent to  $N_1 \geq N_2$ . We can therefore repeat the proof for Theorem 3.5 step by step, except that in (8) we have to use one additional inequality to obtain

$$G_1x = A_1^{-1}N_1x > A_2^{-1}N_1x \geq A_2^{-1}N_2x = G_2x = \rho(G_2)x,$$

and similarly for (9).

To prove Part (ii), assume first that  $A_1 = M_1 - N_1$  is of second type and  $A_2 = M_2 - N_2$  is of first type, so that  $N_1M_1^{-1} \geq O$  and  $M_2^{-1}N_2 \geq O$ . Using (12) we obtain

$$\begin{aligned} M_2^{-1}N_2A_2^{-1} &= M_2^{-1}(M_2 - A_2)A_2^{-1} = A_2^{-1} - M_2^{-1} \\ &\geq A_1^{-1} - M_1^{-1} = M_1^{-1}(M_1 - A_1)A_1^{-1} \\ &= M_1^{-1}N_1A_1^{-1} \\ &= A_1^{-1}N_1M_1^{-1} \geq O. \end{aligned}$$

Now let  $x \geq 0$  and  $y \geq 0$  be two vectors such that

$$N_1M_1^{-1}x = \rho_1x, \quad y^T M_2^{-1}N_2 = y^T \rho_2.$$

These exist by standard Perron–Frobenius theory. Thus

$$\rho_2y^T A_2^{-1}x = y^T M_2^{-1}N_2A_2^{-1}x \geq y^T A_1^{-1}N_1M_1^{-1}x = \rho_1y^T A_1^{-1}x.$$

Since by assumption  $A_1^{-1} > A_2^{-1}$ , and since  $x$  and  $y$  are both nonzero and  $\rho_1 > 0$  we obtain

$$\rho_2y^T A_2^{-1}x > \rho_1y^T A_1^{-1}x.$$

Therefore

$$\rho(M_1^{-1}N_1) < \rho(M_2^{-1}N_2).$$

The case  $M_1^{-1}N_1 \geq O$ ,  $N_2M_2^{-1} \geq O$  can be proved in a similar way.  $\square$

As a first comment, let us note that a special case of the assumption  $A_1^{-1} - A_2^{-1} > O$  arises when  $A_1 \leq A_2$  as we know from Lemma 3.1.

Next, we note that Part (i) of the above theorem generalizes Theorem 3.5 since in the case  $N_1 = N_2$  the assumption (10) is automatically fulfilled (and equality holds there).

Let us further stress the fact that Part (ii) establishes  $\rho(M_1^{-1}N_1)$  as the *smaller* quantity, in contrast to our previous results in Section 3 and to Part (i). So we consider this part as being much more related to the classical single splitting case of Theorem 4.1(iii). Since the conclusions in both parts of Theorem 4.2 are incompatible, the theorem also shows that the respective hypotheses are mutually exclusive. This can essentially also be seen directly: For weak nonnegative splittings of either type we have  $\rho(M_i^{-1}N_i) = \rho(N_iM_i^{-1}) < 1$  so that

$$A_i^{-1} = \sum_{v=0}^{\infty} (M_i^{-1} N_i)^v M_i^{-1} = M_i^{-1} \sum_{v=0}^{\infty} (N_i M_i^{-1})^v,$$

which shows that  $O \leq M_i^{-1} \leq A_i^{-1}$  for  $i = 1, 2$ . Therefore, if we assume (10), i.e.,  $M_2 - M_1 \leq A_2 - A_1$  we obtain

$$\begin{aligned} M_1^{-1} - M_2^{-1} &= M_2^{-1}(M_2 - M_1)M_1^{-1} \\ &\leq A_2^{-1}(M_2 - M_1)A_1^{-1} \\ &= A_2^{-1}(A_2 - A_1)A_1^{-1} \\ &= A_1^{-1} - A_2^{-1}, \end{aligned}$$

which is essentially the opposite of (12).

We now discuss examples which show that Theorem 4.2 no longer holds when certain hypotheses are relaxed. First of all we note that the hypothesis  $A_1^{-1} > O$  cannot be relaxed in Part (i), Example 3.8 representing a counter-example.

Also, the following example shows that without bounds of the kind (10) and (12) we cannot expect comparison results for the spectral radii.

**Example 4.3.** Let

$$\begin{aligned} A_1 &= \begin{pmatrix} 1 & -1 \\ -1 & 2 \end{pmatrix} = \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix} - \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} = M_1 - N_1, \\ A_2 &= \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix} = \begin{pmatrix} 8 & -1 \\ -1 & 2 \end{pmatrix} - \begin{pmatrix} 6 & 0 \\ 0 & 0 \end{pmatrix} = M_2 - N_2, \\ A_3 &= \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix} = \begin{pmatrix} 4 & -1 \\ -1 & 2 \end{pmatrix} - \begin{pmatrix} 2 & 0 \\ 0 & 0 \end{pmatrix} = M_3 - N_3. \end{aligned}$$

All these splittings are  $M$ -splittings and

$$A_1^{-1} = \begin{pmatrix} 2 & 1 \\ 1 & 1 \end{pmatrix} > A_2^{-1} = A_3^{-1} = \frac{1}{3} \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix},$$

and

$$M_1^{-1} = \frac{1}{3} \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix} > M_3^{-1} = \frac{1}{7} \begin{pmatrix} 2 & 1 \\ 1 & 4 \end{pmatrix} > M_2^{-1} = \frac{1}{15} \begin{pmatrix} 2 & 1 \\ 1 & 8 \end{pmatrix}.$$

We have  $M_2 - M_1 \geq O$  but  $M_2 - M_1 \not\leq A_2 - A_1$  as well as  $M_3 - M_1 \geq O$  but  $M_3 - M_1 \not\leq A_3 - A_1$ . So the pairs given by the first and second splittings and by the first and third splittings satisfy all hypotheses of Theorem 4.2, except (10). But we have  $\rho(M_2^{-1}N_2) = 4/5 < \rho(M_1^{-1}N_1) = 2/3 > \rho(M_3^{-1}N_3) = 4/7$ , showing that either inequality between the spectral radii may now occur. Note that we also have  $M_1^{-1} - M_2^{-1} > O$  but  $M_1^{-1} - M_2^{-1} \not\geq A_1^{-1} - A_2^{-1}$  as well as  $M_1^{-1} - M_3^{-1} > O$  but

$M_1^{-1} - M_3^{-1} \not\geq A_1^{-1} - A_3^{-1}$ , i.e., neither pair of splittings satisfies (12). Actually, we have  $M_1^{-1} - M_2^{-1} \leq A_1^{-1} - A_2^{-1}$  and  $M_1^{-1} - M_3^{-1} \leq A_1^{-1} - A_3^{-1}$ . This shows that a modification of Part (ii) with reversed inequalities in (12) and (13) does not hold.

Example 4.3 can be modified slightly to show that Part (ii) of Theorem 4.2 is not empty, i.e., that all hypotheses there can be met. To this purpose, subtract a small positive quantity  $\varepsilon$  from the (1, 2) entry of  $A_3$  and add it at the same position in  $N_3$ . If  $\varepsilon$  is small enough, we will get  $M_3^{-1} - M_2^{-1} \geq A_3^{-1} - A_2^{-1}$  and  $A_3 \preceq A_2$  with  $A_3^{-1} > O$ ,  $A_2^{-1} \geq O$  and both splittings are  $M$ -splittings.

**Remark 4.4.** (i) Interestingly, there exists an obvious, but also new, counterpart of Theorem 4.2 with all strict inequalities replaced by nonstrict ones. (ii) It is possible to formulate a version of Theorem 4.2 where some inequalities between matrices are replaced by less restrictive inequalities between certain matrix vector products. The basic idea can be caught from Theorem 4.5 below, so we do not give details here.

We finish with an additional result where the nonnegativity assumptions on one of the splittings are kept to what we think is a minimum.

**Theorem 4.5.** Assume that  $A_1 = M_1 - N_1$ ,  $A_2 = M_2 - N_2$  are two splittings of nonsingular matrices  $A_1, A_2$  with  $N_1, N_2 \neq O$ . Let  $M_2^{-1}N_2 \geq O$  and assume that there exists an eigenvector  $w \geq 0$  corresponding to  $\rho(M_1^{-1}N_1)$  such that  $0 \leq A_1w$  and

$$A_1w < A_2w. \quad (14)$$

Moreover, let  $O \leq M_1^{-1} \leq M_2^{-1}$ . Then

$$\rho(M_2^{-1}N_2) < \rho(M_1^{-1}N_1). \quad (15)$$

**Proof.** The hypotheses allow us to establish the following chain of (in)equalities:

$$\begin{aligned} \rho(M_1^{-1}N_1)w &= M_1^{-1}N_1w = w - M_1^{-1}A_1w \geq w - M_2^{-1}A_1w \\ &> w - M_2^{-1}A_1A_1^{-1}A_2w = (I - M_2^{-1}A_2)w = M_2^{-1}N_2w. \end{aligned}$$

Since, by assumption,  $M_2^{-1}N_2 \geq O$  this gives  $\rho(M_1^{-1}N_1) < \rho(M_2^{-1}N_2)$  by Lemma 3.2(ii).  $\square$

Theorem 4.5 extends Lemma 2.2 of [10]; cf. also Theorems 3.13 and 3.15 of [8]. The following example shows that this theorem is not contained in our previous ones.

**Example 4.6.** Let

$$A_1 = \begin{pmatrix} -1 & 4/3 \\ 4/3 & -1 \end{pmatrix} = \begin{pmatrix} 0 & 4/3 \\ 4/3 & 0 \end{pmatrix} - \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = M_1 - N_1$$

and

$$A_2 = \begin{pmatrix} 1 & -5/8 \\ -5/8 & 1 \end{pmatrix} = \begin{pmatrix} 1 & -5/9 \\ -5/9 & 1 \end{pmatrix} - \begin{pmatrix} 0 & 5/72 \\ 5/72 & 0 \end{pmatrix} = M_2 - N_2.$$

Then

$$0 \leq M_1^{-1} = \begin{pmatrix} 0 & 3/4 \\ 3/4 & 0 \end{pmatrix} \leq \begin{pmatrix} 81/56 & 45/56 \\ 45/56 & 81/56 \end{pmatrix} = M_2^{-1}.$$

So both splittings are regular splittings. Moreover,  $w = (1, 1)^T$  is a positive eigenvector corresponding to the eigenvalue  $3/4$  (which is the spectral radius) of

$$M_1^{-1}N_1 = \begin{pmatrix} 0 & 3/4 \\ 3/4 & 0 \end{pmatrix}$$

Finally,  $0 \leq A_1 w < A_2 w$ . So all assumptions of Theorem 4.5 are met, while the condition  $M_2 - M_1 \leq A_2 - A_1$  of Theorem 4.2(i) is violated. We have  $\rho(M_2^{-1}N_2) = 5/32$ ,  $\rho(M_1^{-1}N_1) = 3/4$ .

We have some observations concerning the hypotheses of Theorem 4.5. We assume that the eigenvector  $w \geq 0$ , but the proof shows that the hypotheses imply  $w > 0$ . Moreover, since  $0 \leq A_1 w = (M_1 - N_1)w = M_1(I - M_1^{-1}N_1)w = (1 - \rho(M_1^{-1}N_1))M_1 w$  and  $M_1^{-1} \geq 0$ , we get  $\rho(M_1^{-1}N_1) < 1$ ; see [12, Lemma 1] for a similar argument in the special case of a weak nonnegative splitting of first type. By (15), we also have  $\rho(M_2^{-1}N_2) < 1$  which is equivalent to  $A_2^{-1} \geq O$  (see [15, Theorem 3.37]), because the splitting  $A_2 = M_2 - N_2$  is weak nonnegative of first type.

Note that (14) is fulfilled if  $A_2 - A_1 \geq O$  contains no zero rows. Note also that by Theorem 3.4(ii) a vector  $w \geq 0$  such that  $A_1 w \geq 0$  exists if the splitting  $A_1 = M_1 - N_1$  is weak nonnegative of second type.

In Theorem 4.5 the smaller spectral radius corresponds to the larger matrix  $M_2^{-1}$ . But  $M_2^{-1}$  belongs to the splitting of the “smaller” matrix  $A_2^{-1}$ , as the remarks above show. So, as illustrated by Example 4.6, this is yet another situation, different from that of Theorem 4.2. Together with Theorem 3.5 our theorems show the variety of possible results if one compares splittings of *different* matrices.

**Remark 4.7.** Comparison results for splittings of different matrices have previously been used in several ways to study (nonstationary) iterative methods for a (single) system of equations. Typically, one then models the iteration by “macro-iterations” involving splittings of different “macro-matrices”. Examples include the study of multisplittings in [10] (and several subsequent publications) and the paper [6], where the effect of the granularity in the block Jacobi method on its asymptotic rate of convergence is studied. Comparison results on splittings of different matrices appear as an important tool for the analysis in these situations; see, e.g., [[10], Lemma 2.2]. The comparison results of the present paper are an attempt to develop such comparison results in a systematic manner. We believe that they will be useful as

a tool in further investigations on nonstationary iterative processes, including, e.g., inner-outer iterations.

### Acknowledgements

This paper originated from discussions between the authors at Mathematisches Forschungsinstitut Oberwolfach. We are very grateful to all those people who contribute to this very particular *genius loci* which made our work possible and makes Oberwolfach a unique research place in the world. We thank an anonymous referee, whose questions and suggestions led to improvements in our presentation.

### References

- [1] M. Benzi, A. Frommer, R. Nabben, D.B. Szyld, Algebraic theory of multiplicative Schwarz methods, *Numer. Math.* 89 (2001) 605–639.
- [2] A. Berman, R.J. Plemmons, *Nonnegative Matrices in the Mathematical Sciences*, third ed., Academic Press, New York, 1979 (reprinted by SIAM, Philadelphia, 1994).
- [3] J.-J. Climent, C. Perea, Some comparison theorems for weak nonnegative splittings of bounded operators, *Linear Algebra Appl.* 275–276 (1998) 77–106.
- [4] G. Csordas, R.S. Varga, Comparisons of regular splittings of matrices, *Numer. Math.* 44 (1984) 23–35.
- [5] L. Elsner, Comparisons of weak regular splittings and multisplitting methods, *Numer. Math.* 56 (1989) 283–289.
- [6] L. Elsner, M. Neumann, B. Vemmer, The effect of the number of processors on the convergence of the parallel block Jacobi method, *Linear Algebra Appl.* 154–156 (1991) 311–330.
- [7] C.-K. Li, H. Schneider, Applications of Perron–Frobenius theory to population dynamics, *J. Math. Biology* (to appear).
- [8] I. Marek, D.B. Szyld, Comparison theorems for weak splittings of bounded operators, *Numer. Math.* 58 (1990) 387–397.
- [9] I. Marek, D.B. Szyld, Splittings of  $M$ -operators: irreducibility and the index of the iteration operator, *Numer. Funct. Anal. Optim.* 11 (1990) 529–553.
- [10] M. Neumann, R.J. Plemmons, Convergence of parallel multisplitting methods for  $M$ -matrices, *Linear Algebra Appl.* 88–89 (1987) 559–574.
- [11] J.M. Ortega, W.C. Rheinboldt, *Iterative Solution of Nonlinear Equations in Several Variables*, Academic Press, New York, 1970.
- [12] H. Schneider, Positive operators and an inertia theorem, *Numer. Math.* 7 (1965) 11–17.
- [13] H. Schneider, Theorems on  $M$ -splittings of a singular  $M$ -matrix which depend on graph structure, *Linear Algebra Appl.* 58 (1984) 407–424.
- [14] D.B. Szyld, Conditions for the existence of a balanced growth solution for the Leontief Dynamic Input–Output model, *Econometrica* 53 (1985) 1411–1419.
- [15] R.S. Varga, *Matrix Iterative Analysis*, Prentice-Hall, Englewood Cliffs, NJ, 1962 [second edition (revised and expanded) Springer, Berlin, 2000].
- [16] Z.I. Woźnicki, Nonnegative Splitting Theory, *Japan J. Industrial Appl. Math.* 11 (1994) 289–342.
- [17] L. Zeng, Some applications of spectral theory of nonnegative matrices to input–output models, *Linear Algebra Appl.* 336 (2001) 205–218.